



08 a 11 de Outubro de 2018
Instituto Federal Fluminense
Búzios - RJ

ANOMALOUS DIFFUSION THROUGH HOMOPOLAR MEMBRANE: ONE-DIMENSIONAL MODEL

Guilherme Garcia Gimenez¹ – gggimenez@gmail.com

Adélcio C. Oliveira² – adelcio@ufsj.edu.br

¹Discente de Engenharia Mecatrônica e bolsista pela FAPEMIG

²Departamento de Estatística, Física e Matemática

Universidade Federal de São João del-Rei, Campus Alto Paraopeba – Ouro Branco, MG, Brazil

Abstract. We investigate the stationary state solution of the fractional diffusion equation applied to Electrodialysis. This model corresponds to the anomalous diffusion case, that has several applications on Biology, Chemistry and Engineering. With the tools of fractional calculus, we have obtained the analytical solution for the fractional diffusion equation considering a external field. We use the mean field approximation in order to obtain the solution.

Keywords: Anomalous Diffusion, Electrodialysis, Fractional Calculus, Membranes, Mean Field Approximation

1. INTRODUCTION

Electrodialysis is an electrochemical separation process that uses ion-selective membranes as separation agents. Therefore, electrodialysis is a technique in which the separation of substances is accelerated by the application of an adequate electric field. In this way, the ions are transported through these membranes from one solution to another, by means of an electric field as a driving force, without the need of the addition of chemical reagents. Then, cation and anion membranes are organized alternately, in order to form dilute and concentrated flows between the membranes. In a recent work, (Sistat & Pourcelly, 1999) stated the electroneutrality condition can be understood as the fixed and mobile charge have, both, a local null charge. However, this condition is not valid for the entire process of electrodialysis separation and it is not a necessary condition, instead of electroneutrality, (Oliveira *et al.*, 2013; Vieira & Oliveira, 2017) have shown that Mean Field Approximation reproduce better the well known behavior of polarization curve of electrodialysis membrane.

In this work, we investigate the validity of the Mean Field Approximation with the aid of a fractional calculation to analyze the ionic transport through a membrane. This correspond to

the anomalous diffusion case (Almeida & Oliveira, 2018). The anomalous diffusion models have a lot of applications in Biology and Medicine Peng *et al.* (1993); Paiva *et al.* (2009); Hidalgo & Tello (2017), Physics, Astrophysics and Seismology Combe *et al.* (2013); Wegler (2004); Hughes (2010), just to cite a few examples.

2. MATHEMATICAL MODEL FOR ANOMALOUS DIFFUSION

There are many approaches to anomalous diffusion, and also many mathematical definitions of it, see (Almeida & Oliveira, 2018) and references there in. The most widely used is a time dependent stochastic process of random variable x which the mean square displacement is a linear function of time (t), for $t \gg 0$, i.e. $\overline{x^2} \approx Dt^\alpha$, α determines the diffusion type, if $\alpha = 1$ normal diffusion, $\alpha < 1$ sub-diffusion and $\alpha > 1$ super-diffusion.

An important approach to anomalous diffusion is obtained by the porous media equation (Plastino & Plastino, 1995), it is given by:

$$\frac{\partial P}{\partial t} = b^2 \frac{\partial^2 \{P^{2-q}\}}{\partial x^2} \quad (1)$$

where P is the probability density, and b is a constant with dimension of $(Time)(Length)^{-\frac{\alpha}{2}}$. q is related with α by the Tsallis-Bukman scaling law (Tsallis & Bukman, 1996),

$$\alpha = \frac{2}{3-q} \quad (2)$$

the stationary solution of Eq. (1) for an initial delta function is given by:

$$P_q(x) = N_q \left[1 - (1-q) \frac{x^2}{A_q} \right]^{\frac{1}{1-q}} \quad (3)$$

where N_p is a normalizing constant that depends on A_q (Plastino & Plastino, 1995) and A_q is a fit parameter that depends on q . The distribution Eq. (3) is the so called q -Gaussian, that is due to the fact that $\lim_{q \rightarrow 1} P_q$ is the usual Gaussian. It is usual to define the $\beta_q = \frac{1}{A_q}$.

3. ANOMALOUS ELECTRODIALYSIS DIFFUSION MODEL

In this section, we use the transport equation for the separation process in a cell, where a constant electric field (\vec{E}) is applied in the studied system. In a recent work, (Vieira & Oliveira, 2017) suggested that, since the mean field approximation $\vec{E} = \vec{E}_0$, i.e. constant in space directions, was used to support the non-electroneutrality condition, the study should focus on transport in stationary state. To perform the analyzes, we consider that the ions are immersed in a fluid that can be modeled by a diffusion term. In this way, we use the equation and continuity of Nerst-Plank, where the flow of particles can be expressed, in dimensionless form, as:

$$j_k = -d_k \left(\frac{\partial c_k}{\partial x} - z_k c_k E \right) \quad (4)$$

$$\frac{\partial c_k}{\partial t} = -D \frac{\partial j_k}{\partial x} \quad (5)$$

where is an arbitrary scale factor, and all variables and fields are defined in a dimensionless form. Thus, is the dimensionless flux density of the ion , is the diffusion coefficient of ion divided by , is the ion concentration, is the ion valence, is the time, is the position in a opposite direction of the membrane's surface scaled by the membrane thickness. We are interested in the steady state, so we use the fractional equation similar to the stationary equation for the diffusion with Mean Field Approximation, given by:

We are interested in the steady state, so we use the fractional equation similar to the stationary equation for the diffusion with Mean Field Approximation, given by:

$$\frac{\partial j_k}{\partial x} = 0 = -D \left(\frac{\partial^2 c_k}{\partial x^2} - zE_0 \frac{\partial c_k}{\partial x} \right). \quad (6)$$

For a general case using $c_k = u(t)y(x)$, then:

$$\frac{1}{u} \frac{du}{dt} = \frac{D}{y} \left(\frac{d^2 y}{dx^2} - zE_0 \frac{dy}{dx} \right) = -\lambda t \quad (7)$$

where $\lambda > 0$, $\lambda = \frac{w^2}{D}$. Thus, for the spatial part, we will have:

$$\frac{d^2 y}{dx^2} - zE_0 \frac{dy}{dx} + \omega^2 y = 0 \quad (8)$$

Thus, the equivalent fractional equation will be given by:

$$\frac{d^{1+\epsilon} y}{dx^{1+\epsilon}} + k \frac{dy}{dx} + \omega^2 y = 0 \quad (9)$$

where $k = -zE_0$, $\epsilon \in (0,1]$ and $x \in [0, L]$. Therefore, to solve the Eq. (9) we will make use of the Laplace Transform. In this way:

$$y(s) = \mathcal{L}(y(x)) = \int_0^\infty e^{-sx} y(x) dx \quad (10)$$

Then, term by term, we have:

$$\mathcal{L}(\omega^2 y(x)) = \omega^2 y(s) \quad (11)$$

$$\mathcal{L}\left(k \frac{dy}{dx}\right) = k \int_0^\infty e^{-sx} \frac{dy}{dx} dx$$

$$\mathcal{L}\left(k \frac{dy}{dx}\right) = k(ye^{-sx}|_0^\infty + s \int_0^\infty e^{-sx} y dx)$$

$$\mathcal{L}\left(k \frac{dy}{dx}\right) = k(-y(0) + sy(s)) \quad (12)$$

For the fractional term, we have:

$$\mathcal{L}\left(\frac{d^{1+\epsilon}y}{dx^{1+\epsilon}}\right) = \int_0^\infty e^{-sx} \frac{d^{1+\epsilon}y}{dx^{1+\epsilon}} dx \quad (13)$$

defining $u = e^{-sx}$ and $dv = \frac{d^{1+\epsilon}y}{dx^{1+\epsilon}} dx$, we get $v = \frac{d^\epsilon y}{dx^\epsilon}$ and $du = -se^{-sx}$, thus:

$$\begin{aligned} \mathcal{L}\left(\frac{d^{1+\epsilon}y}{dx^{1+\epsilon}}\right) &= \frac{d^\epsilon y}{dx^\epsilon} e^{-sx} \Big|_0^\infty + s \int_0^\infty e^{-sx} \frac{d^\epsilon y}{dx^\epsilon} dx \\ \mathcal{L}\left(\frac{d^{1+\epsilon}y}{dx^{1+\epsilon}}\right) &= -\frac{d^\epsilon y}{dx^\epsilon}(0) + s \int_0^\infty e^{-sx} \frac{d^\epsilon y}{dx^\epsilon} dx \end{aligned} \quad (14)$$

Finally, the integration can be obtained by (Camargo & Oliveira, 2015) as:

$$s \int_0^\infty e^{-sx} \frac{d^\epsilon y}{dx^\epsilon} dx = s^{\epsilon+1}y(s) - s^\epsilon y(0) \quad (15)$$

Therefore, after applying the Laplace Transform in all terms of Eq. (9), we have:

$$\frac{d^{1+\epsilon}y}{dx^{1+\epsilon}} + k \frac{dy}{dx} + \omega^2 y = 0 \Rightarrow s^{\epsilon+1}y(s) - s^\epsilon y(0) - \frac{d^\epsilon y}{dx^\epsilon}(0) + k(-y(0) + sy(s)) + \omega^2 y(s) = 0 \quad (16)$$

Considering $\Omega = \frac{d^\epsilon y}{dx^\epsilon}(0)$ and $\Psi(\Omega) = \Omega + ky(0)$, then we have:

$$\begin{aligned} s^{\epsilon+1}y(s) + s^\epsilon y(0) - \Omega + k(-y(0) + sy(s) + \omega^2 y(s)) &= 0 \\ s^{\epsilon+1}y(s) + ksy(s) + \omega^2 y(s) &= s^\epsilon y(0) + \Omega + ky(0) \\ y(s) &= \frac{s^\epsilon y(0) + \Omega + ky(0)}{s^{\epsilon+1} + ks + \omega^2} \\ y(s) &= \frac{s^\epsilon y(0) + \Psi(\Omega)}{s^{\epsilon+1} + ks + \omega^2}. \end{aligned} \quad (17)$$

The solution can be obtained by inverting Eq. (17), we use the result of (Camargo & Oliveira, 2015), where we have:

$$\mathcal{L}^{-1}\left(\frac{s^{\rho-1}}{s^\alpha + As^\beta + B}\right) = t^{\alpha-\rho} \sum_{r=0}^\infty (-A)^r t^{r(\alpha-\beta)} E_{\alpha, \alpha+1-\rho+(\alpha-\beta)r}^{r+1}(-Bt^\alpha) \quad (18)$$

where $E_{\alpha, \beta}^u(z)$ is the three parameter Mittag-Leffler. Finally, we can write Eq. (17) as:

$$y(s) = y_1(s) + y_2(s) \quad (19)$$

$$y_1(s) = \frac{\Psi(\Omega)}{s^{\epsilon+1} + ks + \omega^2} \quad (20)$$

$$y_2(s) = \frac{s^\epsilon y(0)}{s^{\epsilon+1} + ks + \omega^2} \quad (21)$$

Now, using Eq. (18), we obtain $y_1(s)$ as:

$$\rho = 1, \alpha = \epsilon + 1, A = k, B = \omega^2 \text{ and } \beta = 1, t = x \quad (22)$$

then

$$\mathcal{L}^{-1}(y_1(s)) = y_1(x) = \Psi(\Omega)x^\epsilon \sum_0^\infty (-k)^r x^{r\epsilon} E_{\epsilon+1, \epsilon+1+\epsilon r}^{r+1}(-\omega^2 x^{\epsilon+1}) \quad (23)$$

For $y_2(s)$, we get:

$$\rho = \epsilon + 1, \alpha = \epsilon + 1, A = k, B = \omega^2 \text{ and } \beta = 1, t = x \quad (24)$$

and we have:

$$\mathcal{L}^{-1}(y_2(s)) = y_2(x) = y(0) \sum_0^\infty (-k)^r x^{r\epsilon} E_{\epsilon+1, 1+\epsilon r}^{r+1}(-\omega^2 x^{\epsilon+1}) \quad (25)$$

Finally, the general solution is:

$$y(x) = y_1(x) + y_2(x) \quad (26)$$

3.1 Case $\epsilon = 1$ and $k = 0$

First we consider only the case $\epsilon = 1$. In this way, the Eq. (9) is now:

$$\frac{d^2 y}{dx^2} + k \frac{dy}{dx} + \omega^2 y = 0 \quad (27)$$

then, the solution can be obtained in a form $y = e^{rx}$. The characteristic polynomial is:

$$r^2 + kr + \omega^2 = 0 \quad (28)$$

with $\Delta = k^2 - 4\omega^2$, and with three possible cases, $\Delta > 0$, $\Delta < 0$ and $\Delta = 0$. Then solution is:

$$y_1(x) = \Psi(\Omega)x \sum_{r=0}^\infty (-k)^r x^r E_{2, 2+r}^{r+1}(-(\omega x)^2) \quad (29)$$

$$y_2(x) = y(0) \sum_{r=0}^\infty (-k)^r x^r E_{2, 1+r}^{r+1}(-(\omega x)^2) \quad (30)$$

3.1.1 Case $k = 0$

Now, considering $k = 0$, we have:

$$y_1(x) = \Psi(\Omega)x E_{2, 2}^1(-(\omega x)^2)$$

$$y_1(x) = \frac{\Psi(\Omega)}{\omega} \sin(\omega x) \quad (31)$$

and we have $\frac{\Psi(\Omega)}{\omega} = \frac{dy}{dx}(0)$.

$$y_2(x) = y(0) E_{2, 1}^1(-(\omega x)^2)$$

$$y_2(x) = y(0)E_2(-(\omega x)^2)$$
$$y_2(x) = y(0) \cos(\omega x) \tag{32}$$

4. CONCLUSIONS

As a general remark, we point out that it was possible to generalize the model of (Oliveira et al., 2013) in order to include the anomalous diffusion ingredients. The generalization was carried out with the aid of the Fractional Calculus and its usual techniques. We found a general solution in terms of three parameter Mittag-Leffler function and it was possible to demonstrate that the general solution yields the correct result obtained by (Oliveira et al., 2013) for the non anomalous case. Since membrane are widely used in many applications and naturally appears in biological systems, then we believe that this result can useful to many others researchers.

Acknowledgements

The authors thank the Foundation for Research Support of the State of Minas Gerais (FAPEMIG), through the term APQ-01366-16.

REFERENCES

- Almeida, A. C. L. ; Oliveira, A. C. Diffusion processes modeled with higher order differential terms. FAR EAST JOURNAL OF APPLIED MATHEMATICS, artigo aceito para publicação.
- Baker, R.W., Membrane Technology and Research, Newark, California, 2012.
- Boyce, W. E. & Di Prima, R. C., Equações Diferenciais Elementares e Problemas de Valores de Contorno, ed. Rio de Janeiro: LTC, 2010.
- Camargo, R. F. e Oliveira, E. C., Cálculo Fracionário, Editora Livraria da Física, 2015.
- G. Combe, V. Richefeu, G. Viggiani, S.A. Hall, A. Tengattini and A.P.F. Atman, AIP Conf. Proc. 1542, 453 (2013); doi: 10.1063/1.4811965
- Hidalgo, A., Tello, L., Numerical simulation of a porous medium-type atherosclerosis initiation model, Computers and Fluids (2017), doi:10.1016/j.compfluid.2017.07.019
- Anna L. H. Hughes and Philip J. Armitage, Particle Transport in Evolving Protoplanetary Disks: Implications for Results from Stardust, The Astrophysical Journal, Volume 719, Number 2 (2010).
- Machado, M.B., Avaliação do Processo de Eletrodialise Reversa no Tratamento de Efluentes de Refinaria de Petróleo, Dissertação de Mestrado, Universidade Federal do Rio Grande do Sul, Porto Alegre, 2008.
- Marella, C., Muthukumarappan, K., Metzger, L.E., Application of Membrane Separation Technology for Developing Novel Dairy Food Ingredients. J Food Process Technol 4: 269, 2013.
- Incropera, F. P. and DeWitt, D. P. (2002). Fundamentals of heat and mass transfer. New York: J. Wiley.
- Oliveira, A. C.; Amado, F. D. R. and Moura, R. C. A. Steady state of ion transport in homopolar ionexchange membrane: a theoretical study Journal of the Brazilian Society of Mechanical Sciences and Engineering, (2016),38, 1165-1170.
- L. R. Paiva, C. Binny, S. C. Ferreira Jr., and M. L. Martins, A Multiscale Mathematical Model for Oncolytic Virotherapy , Cancer Res , 69. (3). February 1, (2009) p. 1205.
- Plastino, A. R. and Plastino, A., Physica A 222, 347(1995).
- Peng, C.K. et al. Long-range anticorrelations and non-Gaussian behavior of the heartbeat. Phys. Rev. Lett., v. 70, p. 1343–1346, 1993.
- Rodrigues, M.A.S., Dalla Costa,R.F., Bernardes, A.M., Zoppas Ferreira, J., 2001. Influence of ligand exchange on the treatment of trivalent chromium solutions by electro dialysis, Electrochimica Acta, 47, 2001.
- Scott, K., Handbook of Industrial Membranes, Elsevier Advanced Technology, Oxford, UK, 1995.
- Sistat, P. and Pourcelly, G., (1999). Jour. Elec Chem., 460, 53.

- Strathmann, H., Electrodialysis and related process, in R.D Noble and Stern (ED), Membrane Separations Technology – Principles and Applications, Elsevier, New York, NY, (2005).
- Tsallis, C, and Bukman, D. J., Phys. Rev. E 54, R2197(R) (1996).
- Valero F., Barceló A., Arbós R., Electrodialysis Technology. Theory and Applications, Aigues Ter Llobregat (ATLL), 2007.
- Vieira, I. P.; Oliveira, A. C. Study of Diffusion Applied to Electrodialysis: Three-Dimensional Model in Cylindrical Coordinates. Journal of the Brazilian Society of Mechanical Sciences and Engineering, (2016).
- U. Wegler, Diffusion of seismic waves in a thick layer: Theory and application to Vesuvius volcano, JOURNAL OF GEOPHYSICAL RESEARCH, VOL. 109, B07303, (2004) doi:10.1029/2004JB003048.